

## PUBLICATION INFORMATION

This is the author's version of a work that was accepted for publication in the Tropical Ecology journal. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in <https://doi.org/10.1007/s42965-021-00180-3>

Digital reproduction on this site is provided to CIFOR staff and other researchers who visit this site for research consultation and scholarly purposes. Further distribution and/or any further use of the works from this site is strictly forbidden without the permission of the Tropical Ecology journal.

You may download, copy and distribute this manuscript for non-commercial purposes. Your license is limited by the following restrictions:

1. The integrity of the work and identification of the author, copyright owner and publisher must be preserved in any copy.
2. You must attribute this manuscript in the following format:

This is a manuscript of an article by Georgewill, I., Akani, G.C., Luiselli, L., Petrozzi, F., Dendi, D., Ugbomeh, A.P., Fa, J.E. and Onuegbu, G.C., 2021. **Determining the composition and structure of antelope communities in three study sites within the Niger Delta (Nigeria) based on bushmeat market data.** *Tropical Ecology*. <https://doi.org/10.1007/s42965-021-00180-3>



1       **Composition and structure of the antelope communities at three study**  
2       **areas of the Niger Delta (Nigeria) based on bushmeat market data.**

3

4       Ibiso Georgewill <sup>1</sup>, Godfrey C. Akani <sup>1</sup>, Luca Luiselli <sup>1,2,3</sup>, Fabio Petrozzi <sup>4</sup>, Daniele Dendi  
5                   <sup>1,2,3</sup>, John E. Fa <sup>5,6</sup>, Adaobi P. Ugbomeh <sup>1</sup>, Glorious C. Onuegbu <sup>1</sup>

6

7       <sup>1</sup>Department of Animal & Environmental Biology, Rivers State University of Science and  
8       Technology, P.M.B. 5080, Port Harcourt, Nigeria.

9       <sup>2</sup>Institute for Development, Ecology, Conservation and Cooperation, via G. Tomasi di  
10       Lampedusa 33, I-00144 Rome, Italy

11       <sup>3</sup>Department of Zoology, Faculty of Science, University of Lomé, Lomé, Togo

12       <sup>4</sup>Ecologia Applicata Italia s.r.l., Rome, Italy

13       <sup>5</sup>Division of Biology and Conservation Ecology, School of Science and the Environment,  
14       Manchester Metropolitan University, Manchester M1 5GD, UK.

15       <sup>6</sup>Center for International Forestry Research (CIFOR), Jalan Cifor Rawajaha, Situ Gede,  
16       Bogor Barat, Kota Bogor, Jawa Barat 16115, Indonesia.

17

18       **Running Head:** Niger Delta antelopes

19

20

21

22

23 **Keywords** : Antelopes, bushmeat market, conservation, ethnozoology, Rivers State.

24

25

## INTRODUCTION

26 Nigeria, currently the largest, richest and most densely populated country in Africa, by the  
27 early 1990s was already as densely populated as Western Europe with an average of 130  
28 people per km<sup>2</sup> (Janus and Jaeger, 2011). Since several decades, the rapidly growing  
29 population was placing enormous pressures on the country's natural resource base, and most  
30 of the formerly extensive natural forest and savanna habitats have been degraded or destroyed  
31 by the expansion of agriculture, excessive wood-cutting to supply timber and fuelwood, and  
32 overgrazing of grasslands by livestock (e.g., Osuide, 1990). In addition to widespread  
33 degradation and destruction of natural habitats, wildlife has suffered severely from  
34 uncontrolled hunting for bushmeat (e.g., Martin, 1983; Anadu et al., 1988; Fa et al., 2006). In  
35 southern Nigeria, antelopes have always been purposely targeted among the best target of  
36 hunting for bushmeat because of they are highly valued as tasteful food by local communities  
37 (Luiselli et al., 2019). Thus, their wild populations may have been possibly depleted (e.g.,  
38 Blench, 2007; Luiselli et al., 2015; Petrozzi et al., 2015), despite no quantitative study has  
39 ever been performed in order to evaluate the demographic characteristics and the structure of  
40 Nigerian antelope communities.

41 In order to fill the above-mentioned gap in knowledge, in this paper we analyze the  
42 antelope fauna composition at three distinct study stations in the southern Niger Delta,  
43 characterized by lowland forests and forest-plantation mosaic landscapes. Because of the  
44 difficulty in studying the abundance of these ungulates in the swampy forests, we use data from  
45 three markets to evaluate the relative abundance of the various species and the diversity metrics  
46 of the antelope community. In addition, we also analyze the sex ratio of these populations and  
47 the effect of season on the apparent abundance of the various species.

## 48 MATERIALS AND METHODS

49 The present study was carried out by monitoring three bushmeat markets: Omagwa  
50 (04°59'04"N, 06°55'05"E), Oyigbo (04°53'32"N, 07°10'0"E) and Mbiama (05°03'0"N,  
51 06°27'0"E) in the Rivers State, southern Nigeria (Fig. 1). Rivers State, with over 5 million  
52 inhabitants and more than 630 persons/km<sup>2</sup> density (Rivers State Government, 2019), has  
53 undergone, during the last 30 years, a strong agricultural and industrial expansion that caused  
54 severe fragmentation of the existing forests (Niger Delta Environmental Survey, 1998; Akani,  
55 2008). The study area's climate is characterized by a long rainy season from April through to  
56 the end of September.

57 The three study stations were chosen because they represent localities in which  
58 hunting, alongside traditional agriculture, provide the basis of the local rural population's  
59 economy. The three localities differed in terms of vegetation cover and human population  
60 density (Hansen et al., 2013; Center for International Earth Science Information Network -  
61 CIESIN - Columbia University, 2017); the latter being significantly higher in Mbiama than in  
62 the other localities (Table S1). Local hunters live in bushland and forest patches often <7 km  
63 away from the market. They regularly supply a variety of animal carcasses for their sale.

64 The three bushmeat markets were surveyed during both the dry season (December  
65 2017- March 2018) and the wet season (May 2018- August 2018). Surveying effort was  
66 identical in the three monitored markets: each market was visited (between 7.00-11.00am)  
67 three times per week during eight months (48 daily visits in each season), and all animal  
68 carcasses on sale, including ungulates, were counted on each sampling day. We counted and  
69 inspected the various available carcasses as hunters dropped them with the bushmeat traders.

70 We used contingency table  $\chi^2$  tests to investigate differences among the observed  
71 number of individual animals by sex, season, and market. Saturation curves were built for

72 each market site with 95 % confidence intervals. Bootstrap analysis was applied to generate  
73 upper and lower confidence intervals of all indices, with 9,999 random samples, each with  
74 the same total number of individuals as in each original sample being generated (Harper  
75 1999). Inter-specific differences in the means of a set of morphometric characteristics (Table  
76 S2) were assessed by Student t-test.

77 In order to compare community structure data collected in this study, we used the  
78 following diversity metrics (Magurran, 1988): (a) Species richness, the total number of  
79 species recorded into each habitat type; (b) Dominance:  $D = 1 - \text{Simpson index}$ ; (c) Simpson  
80 index:  $S = 1 - D$ ; (d) Shannon-Wiener  $H'$  index (Shannon & Weaver, 1963; (e) Evenness,  
81 calculated using Pielou's formula (Magurran, 1988); (f) Chao 1, the number of species  
82 predicted to be present at each study area given the sample observed (Hughes et al. 2001;  
83 Chodak et al. 2013). We calculated the 95% upper and lower confidence intervals using  
84 10,000 bootstraps. Alpha level was set at  $p = 0.05$ . Past 3.0 software was used to calculate the  
85 various diversity indices.

## 86 RESULTS

87 During the research work, a total of 202 Antelopes was counted (Table 1) within the  
88 sampling duration. The frequency of antelope carcasses differed significantly by market site  
89 ( $\chi^2 = 36.6$ ,  $df = 2$ ,  $P < 0.0001$ ), with most animals being traded in Omagwa ( $n = 126$  antelope  
90 carcasses), followed by Oyigbo ( $n = 47$ ) and Mbiama ( $n = 29$ ). In all markets, the same three  
91 species were recorded: *Tragelaphus scriptus* ( $n = 24$ ), *Philantomba walteri* ( $n = 141$ ), and  
92 *Neotragus batesi* ( $n = 35$ ) (Figure 1). Sex ratio was even in the three species: *Tragelaphus*  
93 *scriptus* (1.6 male : 1 female;  $\chi^2 = 0.7$ ,  $df = 1$ ,  $P = 0.402$ ), *Philantomba walteri* (0.98 male : 1  
94 female;  $\chi^2 = 0.003$ ,  $df = 1$ ,  $P = 0.953$ ), and *Neotragus batesi* (0.94 male : 1 female;  $\chi^2 = 0.014$ ,  
95  $df = 2$ ,  $P = 0.905$ ).

96 64.3% of the antelope carcasses were traded during the wet season (inter-seasonal  
97 differences:  $\chi^2 = 8.5$ ,  $df = 1$ ,  $P < 0.01$ ). The inter-seasonal differences were not significant in  
98 *Tragelaphus scriptus* ( $\chi^2 = 0.31$ ,  $df = 1$ ,  $P = 0.578$ ) and *Neotragus batesi* ( $\chi^2 = 3.32$ ,  $df = 1$ ,  $P$   
99  $= 0.068$ ), whereas they were statistically significant for *Philantomba walteri* ( $\chi^2 = 5.6$ ,  $df = 1$ ,  
100  $P < 0.01$ ).

101 In terms of diversity metrics, Omagwa appeared ecologically better than the other  
102 sites, with Mbiama being intermediate and Oyigbo being more depleted: indeed, despite the  
103 taxonomical composition of the species was identical across sites, the evenness and diversity  
104 indices were highest in Omagwa and lowest in Oyigbo, whereas the opposite was true for the  
105 dominance index (Table 2). Interestingly, saturation curves also revealed that, whereas in  
106 Omagwa no other antelope species can be expected, the same was not true for the other two  
107 sites where the plateau of the curve was not reached (Figure 2).

108 The morphometric characteristics of the different species of antelopes across stations  
109 are given in Table S2, showing that, as expected, *Tragelaphus scriptus* carcasses were  
110 significantly larger than the other two species in all body measures (in all cases,  $P < 0.0001$  at  
111 Student t-tests). The market value depended directly on the relative size of the carcasses:  
112 thus, *Tragelaphus scriptus* was sold at higher prices than the other two species (Table 3).

## 113 **DISCUSSION**

114 Recent literature has showed that there is considerable confusion concerning the antelope  
115 species in the Niger Delta: for example, for duikers (genera *Cephalophus* and *Philantomba*),  
116 only one of the six species cited in the literature were demonstrated as definitely present in  
117 the Niger Delta region, and, overall, only six antelope species have been recorded out of  
118 which only five were recorded more than once (Petrozzi et al., 2015). Thus, the species  
119 diversity of Niger Delta antelopes is by far less than historically reported (e.g., Happold,

120 1987; Powell, 1993; Angelici et al., 1999; Blench, 2007), also because it is likely that some  
121 species were wrongly reported for the general area (Petrozzi et al., 2015; Luiselli et al., 2015,  
122 2019a). Therefore, the reduced species richness ( $n = 3$ ) observed in the three study stations is  
123 not surprising, whereas the total number of carcasses ( $n = 202$ ) was low if we consider (i) the  
124 considerable field effort, (ii) the appreciation for antelope meat by local communities  
125 (Luiselli et al., 2019b) and (iii) the fact that three distinct localities were monitored.  
126 *Philantomba walteri* was the dominant species at all the three study stations as it constitutes  
127 about 70% of the total antelope carcasses recorded. This data fully mirrors data presented by  
128 Petrozzi et al. (2015). At another site in a forested area of the central Niger Delta,  
129 *Philantomba walteri* was also the most abundant species, but also *Tragelaphus scriptus* and  
130 *Tragelaphus spekei* (not seen in the present study) had practically the same abundance,  
131 whereas *Neotragus batesi* and *Hyaemoschus aquaticus* (also not recorded during the present  
132 study) appeared less abundant (Akani et al., 2015). Comparatively, it seems that the antelope  
133 species richness is still higher in the forests of the western and central side of the Niger Delta  
134 than in the eastern side of the deltaic axis, with only one species (*Philantomba walteri*) being  
135 still widespread and abundant. However, the fact that a relatively low number of antelope  
136 carcasses has been observed along the study period (much lower than the number of carcasses  
137 of small carnivores for instance, see Onuegbu et al., submitted) suggests that the ungulate  
138 fauna is already very depleted in the eastern Niger Delta region, as also observed in other  
139 African areas with heavy hunting pressure (Fa et al., 1995; Fischer and Linsenmair, 2001;  
140 Grande-Vega et al., 2016; Hema et al., 2017).

141 Sex ratio of our observed samples was even for all species. Literature data suggests  
142 that sex ratio may vary considerably in *Tragelaphus scriptus* from area to area with some  
143 populations having even sex ratios and others having female-skewed ratios (e.g., Waser,  
144 1975; Alsopp, 1978; Yazezew et al., 2011) whereas nothing is known on *Philantomba walteri*

145 and *Neotragus batesi*. Thus, although preliminary, our study gives the first-of-ever data on  
 146 the adult sex ratio of a reasonable sample for these two ecologically nearly unknown antelope  
 147 species.

148 Our data also suggest that antelopes (in particular *Philantomba walteri*) are hunted  
 149 more intensely by wet season. These data mirror the same patterns observed with small  
 150 carnivores sold in bushmeat markets (Onuegbu et al., submitted), and are in agreement with  
 151 information provided by hunters (n = 66) that reported hunting to be more productive at wet  
 152 season than at dry season. According to them, the sound from dry grasses/vegetation during  
 153 dry season provided an easy escape route for the antelopes thereby reducing their catches  
 154 compared to rainy season. According to most of the interviewed hunters, *Philantomba walteri*  
 155 and *Neotragus batesi* usually occupy low successional disturbed habitats, unlike *Tragelaphus*  
 156 *scriptus* that prefers undisturbed habitats.

## 157 REFERENCES

- 158 Akani, G.C., Amadi, N., Eniang, E.A, Luiselli, L. & Petrozzi, F. (2015). Are mammal  
 159 communities occurring at a regional scale reliably represented in “hub” bushmeat  
 160 markets? A case study with Bayelsa State. (Niger Delta, Nigeria). *Folia Zoologica*,  
 161 64, 79-86.
- 162 Allsopp, R. (1978). Social biology of bushbuck (*Tragelaphus scriptus* Pallas 1776) in the  
 163 Nairobi National Park, Kenya. *African Journal of Ecology*, 16(3), 153-165.
- 164 Angelici FM, Grimod I, Politano E. 1999. Mammals of the eastern Niger Delta (Rivers and  
 165 Bayelsa States, Nigeria): an environment affected by a gas-pipeline. *Folia Zoologica*.  
 166 48:249– 264.



- 167 Anadu, P. A., Elamah, P. O., & Oates, J. F. (1988). The bushmeat trade in southwestern  
 168 Nigeria: a case study. *Human Ecology*, 16(2), 199-208.
- 169 Blench R. 2007. Mammals of the Niger Delta, Nigeria. Privately printed p. 64.
- 170 Center for International Earth Science Information Network - CIESIN - Columbia University.  
 171 (2017). Gridded Population of the World, Version 4 (GPWv4): Population Density,  
 172 Revision 10. Palisades, NY: NASA Socioeconomic Data and Applications Center  
 173 (SEDAC). <https://doi.org/10.7927/H4DZ068D>. Accessed 28/08/2018
- 174 Chodak, M., Gołębiewski, M., Morawska-Płoskonka, J., Kuduk, K., and Niklińska, M.  
 175 (2013). Diversity of microorganisms from forest soils differently polluted with heavy  
 176 metals. *Applied Soil Ecology* 64:7-14.
- 177 Fa, J.E., Yuste, J., Perez Del Val, J.&Castroviejo, J. (1995) Impact of market hunting on  
 178 mammal species in Equatorial Guinea. *Conservation Biology*, 9, 1107–1115.
- 179 Fa, J. E., Seymour, S., Dupain, J. E. F., Amin, R., Albrechtsen, L., & Macdonald, D. (2006).  
 180 Getting to grips with the magnitude of exploitation: bushmeat in the Cross–Sanaga  
 181 rivers region, Nigeria and Cameroon. *Biological Conservation*, 129(4), 497-510.
- 182 Fischer, F. & Linsenmair, K.E. (2001) Decreases in ungulate population densities. Examples  
 183 from the Comoé National Park, Ivory Coast. *Biological Conservation*, 101, 131–135.
- 184 Grande-Vega, M., Farfan, M.A., Ondo, A. & Fa, J.E. (2016) Decline in hunter offtake of blue  
 185 duikers in Bioko Island, Equatorial Guinea. *African Journal of Ecology*, 54, 49–58.
- 186 Hansen, M. C., P. V. Potapov, R. Moore, M. Hancher, S. A. Turubanova, A. Tyukavina, D.  
 187 Thau, S. V. Stehman, S. J. Goetz, T. R. Loveland, A. Kommareddy, A. Egorov, L.  
 188 Chini, C. O. Justice, and J. R. G. Townshend. (2013). “High-Resolution Global Maps of  
 189 21st-Century Forest Cover Change.” *Science* 342 (15 November): 850–53. Data  
 190 available on-line from: [http://earthenginepartners.appspot.com/science-2013-global-](http://earthenginepartners.appspot.com/science-2013-global-forest)  
 191 forest.

- 192 Happold DCD. 1987. The mammals of Nigeria. Oxford: Clarendon Press
- 193 Harper D.A.T. (ed.) 1999: Numerical palaeobiology. John Wiley & Sons, New York.
- 194 Hema, E. M., Ouattara, V. A. L. Y., Parfait, G., Di Vittorio, M., Sirima, D., Dendi, D., ... &  
195 Luiselli, L. (2017). Bushmeat consumption in the West African Sahel of Burkina Faso,  
196 and the decline of some consumed species. *Oryx*, doi:10.1017/S0030605316001721.
- 197 Hughes, J. B., Hellmann, J. J., Ricketts, T. H., and Bohannan, B. J. (2001). Counting the  
198 uncountable: statistical approaches to estimating microbial diversity. *Appl. Environ.*  
199 *Microbiol.* 67:4399-4406.
- 200 Janus, A., and Jaeger, R. (2011). Country Profile Nigeria. *2011 Country Profiles*, 1–8.  
201 <https://doi.org/10.1111/2041-9066.12008>
- 202 Luiselli, L., Amori, G., Akani, G.C. & Eniang, E.A. (2015). Ecological diversity, community  
203 structure and conservation of Niger Delta mammals. *Biodiversity and Conservation*, 24,  
204 11, 29-34.
- 205 Luiselli, L., Dendi, D., Eniang, E. A., Fakae, B. B., Akani, G. C., & Fa, J. E. (2019a). State of  
206 knowledge of research in the Guinean forests of West Africa region. *Acta Oecologica*  
207 94: 3-11.
- 208 Luiselli, L., Hema, E. M., Segniagbeto, G. H., Ouattara, V., Eniang, E. A., Di Vittorio, M., ...  
209 & Sirima, D. (2019b). Understanding the influence of non-wealth factors in determining  
210 bushmeat consumption: results from four West African countries. *Acta Oecologica* 94:  
211 47-56.
- 212 Magurran, A.E. (1988). Ecological diversity and its measurement. Princeton, New Jersey:  
213 Princeton University Press.
- 214 Martin, G. H. G. (1983). Bushmeat in Nigeria as a natural resource with environmental  
215 implications. *Environmental Conservation*, 10(2), 125-132.
- 216 Osuide, S. O. (1990). Environmental pollution in Nigeria. *Habitat International*, 14(1), 5–15.

- 217 [https://doi.org/10.1016/0197-3975\(90\)90014-R](https://doi.org/10.1016/0197-3975(90)90014-R)
- 218 Petrozzi, F., Akani, G.C., Amadi, N., Eniang, E.A., Gippoliti, S., Luiselli, L., 2015. Surveys  
219 of mammal communities in a system of five forest reserves suggest an ongoing biotic  
220 homogenization process for the Niger Delta (Nigeria). *Trop. Zool.* 28, 95-113.
- 221 Powell, C. B. (1993): Sites and Species of Conservation Interest in the Central Axis of the  
222 Niger Delta. Abuja, Natural Resource Conservation Council.
- 223 Rivers State Government (2019). Available at: <http://www.riversstate.gov.ng> [Accessed on  
224 17 May 2019].
- 225 Waser PM (1975) Spatial association and social interactions in a solitary ungulate: the  
226 bushbuck *Tragelaphus scriptus* (Pallas). *Z Tierpsychol* 37:24–36.
- 227 Yazezew, D., Mamo, Y., & Bekele, A. (2011). Population Ecology of Menelik’s Bushbuck  
228 (*Tragelaphus scriptus meneliki*, Neumann 1902) from Denkoro Forest Proposed  
229 National Park, Northern Ethiopia. *International Journal of Ecology and Environmental*  
230 *Sciences*, 37(1), 1-13.
- 231

232 **Table 1.** Synopsis of the antelope data collected at the three study stations during the research  
 233 period.

234

---

Species	Station 1 (Omagwa)	Station 2 (Oyigbo)	Station 3 (Mbiama)
<i>Tragelaphus scriptus</i>	24	1	1
<i>Philantomba walteri</i>	82	38	21
<i>Neotragus batesi</i>	20	8	7
TOTAL	126	47	29

235

236

237 **Table 2.** Estimates of diversity metrics for antelope assemblages (as indicated by bushmeat  
 238 market surveys) in southern Nigeria, after 10,000 bootstraps. Lower = lower 95% confidence  
 239 interval; upper = upper 95% confidence interval.

240

	Omagw a	Lowe r	Uppe r	Oyigb o	Lowe r	Uppe r	Mbiam a	Lowe r	Uppe r
Taxa_S	3	3	3	3	3	3	3	3	3
Individuals	126			47			29		
Dominance	0.49	0.42	0.57	0.68	0.54	0.81	0.58	0.45	0.76
Simpson	0.52	0.43	0.58	0.32	0.19	0.46	0.42	0.24	0.55
Shannon	0.89	0.76	0.98	0.56	0.39	0.76	0.69	0.48	0.91
Evenness	0.81	0.71	0.89	0.58	0.49	0.71	0.67	0.54	0.83
Chao-1	3	3	3	3	3	3	3	3	3

241

242

243 **Table 3.** Market value of antelope carcasses by station and by species.

244

245

Species	Station	Price Range (Naira)	Mean Price (Naira)
<i>Philantomba walteri</i>	Omagwa	6000-11000	7500
	Oyigbo	7000-9000	8000
	Mbiana	6800-10000	7850
<i>Neotragus batesi</i>	Omagwa	5000-8000	6500
	Oyigbo	4500-8000	6000
	Mbiana	4500-8000	6000
<i>Tragelaphus scriptus</i>	Omagwa	14000-25000	10500
	Oyigbo	8500-13500	10000
	Mbiana	8000-12000	9500

246


247

248

249

250 **Figure 1:** Map of Rivers State in southern Nigeria, showing the three sample stations, and the  
251 three study species: *Neotragus batesi* (b), *Philantomba walteri* (c) and *Tragelaphus scriptus*  
252 (d)

253




254

255

256 **Figure 2.** (A) Saturation curves (with 95 % confidence intervals after 9999 bootstraps) and  
 257 (B) Diversity profiles (95 % confidence, after 9999 bootstraps), for the community diversity  
 258 of antelopes at the three study stations


259

260 (A)



261

262 (B)



263



264 **ONLINE SUPPLEMENTAL MATERIAL**

265

266 **Table S1:** GIS-based estimates of the dominant tree cover (in terms of % of occupied land)  
 267 and of the human population density, for a 7-km-radius buffer along the three surveyed  
 268 market sites and another area of the Niger Delta (Swali) used for literature comparison. Data  
 269 from Hansen et al. (2013) and Center for International Earth Science Information Network -  
 270 CIESIN - Columbia University (2017).

271

Surveyed locality	% of dominant tree cover	human population density (per km <sup>2</sup> )
Swali	56	638.2
Mbiama	55	760.1
Oyigbo	16	380.8
Omagwa	29	371.6

272

**Table S2.** Summary of the morphometric characteristics of the different species of antelopes across stations (mean±SD)

Species	Location	Sex	Weight (kg)	Total Length (cm)	Standard length (cm)	Hind foot (cm)	Forefoot (cm)	Ear (cm)	Body depth (cm)	Tail (cm)	Head (cm)	Neck (cm)
<i>Philantomba walteri</i>	Mbiama	F	6.1±2.31	27.75±2.73	23.06±2.48	11.56±2.6	10.19±2.58	2.89±0.33	10.25±1.16	4.88±1.6	5.31±0.8	3.49±0.25
		M	6.83±2	26.33±6.04	22±5.48	12.61±2.48	11.06±1.99	3.16±0.33	10.94±1.21	5.39±1.87	5.83±1.03	3.63±0.29
	Oyigbo	F	6.77±2.01	28±2.7	23.38±2.38	12.54±3.31	10.29±2.39	3.04±0.36	10.54±1.23	5.46±1.57	5.63±0.91	3.58±0.28
		M	6.48±1.96	26.57±4.7	22.23±4.29	11.97±2.54	10.53±2.18	3.05±0.34	10.6±1.27	5.23±1.73	5.57±0.96	3.59±0.27
	Omagwa	F	6.78±2.05	28.41±2.36	23.64±1.99	11.83±2.55	10.33±2.23	3.06±0.35	10.55±1.09	5.47±1.49	5.68±0.86	3.61±0.28
		M	6.19±1.78	24.94±5.29	20.75±4.83	11.71±2.17	10.52±2	3.05±0.34	10.52±1.13	4.91±1.72	5.42±0.78	3.66±0.37
<i>Neotragus batesi</i>	Mbiama	F	5.03±0.45	16.83±2.02	13.67±2.08	11.83±2.02	10.83±2.02	4.03±0.45	10.67±1.26	3.33±0.58	4.83±0.21	3.9±0.1
		M	5.14±0.31	17.17±1.61	14±1.73	12.17±1.61	11.17±1.61	4.1±0.36	10.83±1.04	3.33±0.58	4.9±0.1	3.9±0.1
	Oyigbo	F	5.01±0.46	16.67±2.08	13.67±2.08	11.67±2.08	10.67±2.08	3.97±0.47	10.5±1.32	3.33±0.58	4.77±0.15	3.83±0.06
		M	5.21±0.41	17.5±2.12	14.5±2.12	12.5±2.12	11.5±2.12	4.15±0.49	11±1.41	3.5±0.71	4.85±0.07	3.85±0.07
	Omagwa	F	4.89±0.15	16.07±0.53	12.86±0.38	11.07±0.53	10.07±0.53	3.84±0.16	10.14±0.38	3.07±0.19	4.86±0.15	3.99±0.25
		M	5.35±0.68	17.64±2.32	14.71±2.63	12.64±2.32	11.64±2.32	4.26±0.61	11±1.29	3.57±0.73	4.89±0.17	4±0.26
<i>Tragelaphus scriptus</i>	Mbiama	F	40±0	36±0	30±0	18±0	15±0	3±0	16.5±0	7±0	6.5±0	6.5±0
	Oyigbo	M	58±0	47±0	39±0	22±0	19.5±0	4.5±0	18±0	9±0	8±0	7±0
	Omagwa	F	46.5±3.11	44.775±2.05	35.625±1.49	19.375±1.11	17.75±1.19	3.833333±0.29	18.125±1.65	8±1.22	7.125±0.75	6.75±0.5
		M	52.91±5.22	47.07±3.42	37.5±3.16	20.94±1.86	18.72±1.66	4.81±1.03	15.44±1.67	9.17±0.56	7.94±0.88	7.39±0.74

