

Building a framework towards climate-smart agriculture in the Yangambi landscape, Democratic Republic of Congo (DRC)

Climate-smart
agriculture

Lisette Mangaza

*Faculty of Sciences, University of Goma, Goma, Congo
(the Democratic Republic Of The)*

Denis Jean Sonwa

Center for International Forestry Research (CIFOR), Yaoundé, Cameroon

Germain Batsi and Jérôme Ebuy

*Faculty of Renewable Natural Resources Management, University of Kisangani,
Kisangani, Congo (Democratic Republic of Congo), and*

Jean-Marie Kahindo

*Faculty of Sciences, University of Kisangani, Kisangani,
Congo (Democratic Republic of Congo)*

Received 7 August 2020
Revised 16 January 2021
12 February 2021
18 February 2021
15 March 2021
Accepted 19 March 2021

Abstract

Purpose – This paper aims to produce a framework for climate-smart agriculture (CSA) in the Yangambi landscape, Democratic Republic of the Congo (DRC). This would enable the authors to identify agricultural practices, assess vulnerability to climate change, identify options for improving agricultural systems from a climate change mitigation and adaptation perspective and finally provide climate-smart agricultural options.

Design/methodology/approach – The study used household survey methods of data collection. The data were collected using a structured questionnaire survey by interviewing 250 farm households, subdivided

© Lisette Mangaza, Denis Jean Sonwa, Germain Batsi, Jérôme Ebuy and Jean-Marie Kahindo. Published by Emerald Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licenses/by/4.0/legalcode>

The authors would like to thank the European Union and the Centre for International Forestry Research (CIFOR) for funding, through the Training and Research in Tshopo (FORETS) project, this research carried out as part of master's degree work but also as part of the CIFOR Global Comparative Study for achieving REDD+ (GCS-REDD+) project funded by the Norwegian Agency for Development Cooperation (NORAD). The authors would also like to thank the R&SD (Resources and Synergies Development) design office for its support in logistics. The authors would like to thank all farmers in the Yangambi landscape for their support during the data collection period.

Author contributions: Denis Sonwa conceptualized and supervised the study with Jean-Marie Kahindo and Jérôme Ebuy; Lisette Mangaza collected the data with the assistance of Germain Batsi. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by EU (Contrat FED/2016/381–145) and NORAD Grant agreement code No: NOR114.



using three axes of the Yangambi landscape. Fisher's exact test was used to determine relationships between two or more variables.

Findings – Results of the survey revealed that the vast majority (98%) of respondents perceived changes in temperature, rainfall and weather patterns. Reduction of crop yields and the emergence of new weed species and new crop pests are the main impacts on agricultural activities. Although 87.6% of respondents have no means of adaptation and resilience, some of them use crops rotation, fallow practice, fertilizers and bio-pesticides. A framework for CSA is proposed for the Yangambi landscape.

Practical implications – Policies and strategies to promote CSA in the study area should take into account local farmers' perceptions of climate change and consider first the adequacy of CSA practices for the specific conditions of the target area before its promotion. This study is thus useful for many REDD+ initiatives that are currently being promoted in DRC and particularly in the Tshopo Province.

Originality/value – This study is one of the first studies to focus on CSA in the Yangambi landscape, DRC. It assists the use of agriculture as a response to reducing deforestation while at the same time lowering agriculture's carbon footprint and promoting a resilient and more productive farming system.

Keywords Climate change, Climate-smart agriculture, Congo Basin, DRC (Democratic Republic of the Congo), Farmers' perceptions, Yangambi

Paper type Research paper

1. Introduction

Climate change is already hampering agricultural activities. According to the Intergovernmental Panel on Climate Change (IPCC), climate change affects crop production, with negative effects more common than positive (IPCC, 2014). Africa is the most vulnerable continent because of its heavy dependence on rain-fed agriculture (IPCC, 2007), widespread poverty, low adaptive capacity and lack of investment in mitigation and resilience-building systems (Bele *et al.*, 2010; Williams *et al.*, 2015).

Over the centuries, farmers have learned to adapt to climate variability, but given the pace and intensity of current and future climate change, their actions are no longer sufficient (Bele *et al.*, 2010). Rural areas are the most vulnerable (IPCC, 2007; FAO, 2016).

However, this vulnerability is greater in Congo Basin countries such as the Democratic Republic of the Congo (DRC) (Sonwa *et al.*, 2012; Ministry of Foreign Affairs, 2018), where about 70% of the population live in rural areas and depend on rain-fed slash-and-burn agriculture to live (Ulimwengu and Kibonge, 2016). In the Yangambi landscape, agriculture is the main activity and accounts for between 70% and 85% of the income of three-quarters of households. Economic opportunities are scarce and more than 80% of households live below the poverty line (CIFOR, 2018a).

Moreover, despite the fact that agriculture is considered as the most vulnerable sector to climate change at both local and national levels (MECNT and UNDP, 2009; Ulimwengu and Kibonge, 2016); several studies have shown that it is a major source of greenhouse gas (GHG) emissions (Pan *et al.*, 2011; Arneth *et al.*, 2019), through the process of deforestation (Potapov *et al.*, 2012; Hufkens *et al.*, 2020), thus contributing to climate change (Kipalu and Mukungu, 2012; AMCEN, 2014). Slash-and-burn agriculture is the main direct cause of deforestation in DRC (MECNT, 2012; Molinario *et al.*, 2015), particularly in the Yangambi landscape (Hufkens *et al.*, 2020). The search for fertile land pushes the population to clear increasingly more forest areas to meet their ever-increasing needs and to cope with greater demographic pressure (Alongo *et al.*, 2013; Kyale *et al.*, 2019).

From the above, it can be seen that there are two major agricultural challenges in the Yangambi landscape:

- (1) combating climate change by introducing agricultural practices that aim to reduce GHG emissions, while adapting to climate change (Seguin, 2010); and

-
- (2) increasing agricultural production in order to ensure food security while also preserving forests.

Thus, to alleviate the agricultural challenge in a climate change context, the Food and Agriculture Organization (FAO) of the United Nations suggested climate-smart agriculture (CSA); this is an approach based on three pillars – food security, adaptation and mitigation (Delvaux, 2016). Its agricultural practices increase productivity and resilience in a sustainable manner, reduce GHG emissions intensity, curb deforestation and improve the health of soils, landscapes and forests (FAO, 2016).

CSA is not a prescribed practice or a specific technology that can be universally applied. It requires adoption of an integrated approach that takes into account specific local conditions (FAO, 2014), including site-specific assessments of the social, economic and environmental conditions to identify appropriate agricultural technologies and practices (Williams *et al.*, 2015).

So, within this framework, we conducted this study, of which the main objective is to establish a framework towards CSA in the Yangambi landscape. Specifically, the study aims to:

- identify current agricultural practices and their constraints in terms of yield;
- assess vulnerability to climate change and identify adaptation options;
- identify options for improving the performance of agricultural systems from a climate change mitigation and adaptation perspective; and
- finally, provide climate-smart agricultural options for the Yangambi landscape.

2. Materials and methods

2.1 Study area

This work was carried out in the Yangambi landscape, located in Tshopo Province, in Northeastern DRC (Figure 1). The region corresponds to an area located between 24°16'95" and 25°08'48" E longitude; 0°38'77" and 1°10'20" N latitude. The climate of Yangambi is continental equatorial, belonging to the Af type of the Köppen classification (Van Wambeke and Evrard, 1957). The annual precipitation varies between 1600 and 2200 mm (Mohyont and Demarée, 2006), with an annual average of 1820 mm (Likoko *et al.*, 2019). The annual temperature varies from 24.2°C to 25.8°C, with an annual average of 24.98°C (Kombele, 2004; Likoko *et al.*, 2019). The main activity in the Yangambi landscape is slash-and-burn subsistence agriculture. Cassava, maize, rice, groundnut and plantain are the main crops (CIFOR, 2018a).

2.2 Methods

2.2.1 Research design. The study was carried out in nine sites distributed along three axes in the Yangambi landscape: Yangambi–Ngazi road axis, Yangambi center and Kisangani–Yangambi road axis (Table 1). The Yangambi center or the city of Yangambi is an area housing the National Institute for Agronomic Studies and Research (INERA)/Yangambi, whose activities are mainly influenced by INERA. The Kisangani–Yangambi axis is along the road linking Yangambi to the city of Kisangani, capital city of the province. Activities are more influenced by the city of Kisangani than by the city of Yangambi. The Yangambi–Ngazi axis is a rural area not connected directly to a large city.

2.2.2 Data collection. The study used household survey methods of data collection. The data were collected using a structured questionnaire survey. Information collected includes,




Figure 1.
Localization of the study area

Axes	Sites	No. of households surveyed	No. of households surveyed per axis
Yangambi center	Bangala	35	140
	Likango	35	
	Lumumba	35	
	Lusambila	35	
	Yalinga	25	
Yangambi–Ngazi axis	Yanguma	20	45
	Yakako	20	
Kisangani–Yangambi axis	Yalungu	20	65
	Yaselia	25	
	Yakako1	20	
All three axes		250	250

Table 1.
Study sample

household characteristics, agricultural practices and constraints, and farmers’ perceptions of climate change. The survey was administered from 23 May to 12 June 2018. Qualitative and quantitative data were collected from farmers having at least five years of seniority in the site, being head of the household and having a field. A total of 250 households were surveyed. Secondary information was collected through focus group discussions. These consisted of about 10 people at each site before the household survey was administered, to gain a general idea about the region.

2.2.3 Data analysis. Fisher’s exact test was used to determine if there was a significant relationship between two or more variables. It was mainly used to establish a relationship between different farmers’ perceptions of climate change and belonging to an axis. The acceptable error for the statistical analyses was 5%. When the *p*-value was less than the

significance level of 5%, this meant that there was a significant relationship between variables. These statistical analyses were performed using RStudio software version 3.3 (Barnier *et al.*, 2020).

3. Results

3.1 Agricultural practices in the Yangambi landscape

Households own on average 1.67 ± 0.79 fields of 0.75 ± 0.52 hectares in the Yangambi landscape (data not shown). Most of them are established on fallow land (85.2%), whereas others are in mature forest (11.6%) or are planted directly after the harvest of another crop (3.2%).

3.1.1 Presence of trees in the fields. Some trees are left in fields by farmers during the establishment of new fields, whereas others grow spontaneously between crops and are retained during maintenance work. The most reported trees are *Petersianthus macrocarpus* (P. Beauv.) Liben (18.4% of respondents), *Persea americana* Mill. (6.8% of respondents), *Albizia ferruginea* (Guill. and Perr.) Benth (4.8% of respondents), *Pychmantis angolensis* (Welw.) Warb. (4.4% of respondents), *Annonidium mannii* (Oliv.) Engl. and Diels (4.0% of respondents) and *Dacryodes edulis* (G. Don) H. J. Lam (2.4% of respondents). Trees are retained in the fields for different reasons. Thus, *Petersianthus macrocarpus* (called “boso” in the local language) is kept for the edible caterpillars it bears, whereas *Persea americana* (avocado tree) and *Dacryodes edulis* (African pear) are kept for their edible fruits.

Apart from trees already in the fields, there are some trees species that respondents would prefer in their fields and would be willing to plant them. The most reported by respondents are *Petersianthus macrocarpus* (40.4% of respondents), *Persea americana* (24.4% of respondents), *Dacryodes edulis* (15.6% of respondents) and *Erythrophleum suaveolens* (Guill. and Perr.) Brenan (13.2% of respondents). *Erythrophleum suaveolens* is preferred for its medicinal properties, whereas other trees are preferred for reasons mentioned above.

3.1.2 Main crops in the Yangambi landscape. The main crops in the Yangambi landscape are cassava (*Manihot esculenta* Crantz, 62% of respondents), rice (*Oryza sativa* L., 22.4% of respondents), maize (*Zea mays* L., 6% of respondents), groundnut (*Arachis hypogea* L., 5.2% of respondents), cowpea (*Vigna unguiculata* (L.) Walp., 2.4% of respondents) and plantain and table banana (*Musa spp.* L., 2.0% of respondents). Apart from cassava, which is the main crop throughout the entire Yangambi landscape, there is a significant difference in the choice of other crops by axes ($p = 0.004$). Thus, rice predominates along the Yangambi–Ngazi axis, whereas cowpea predominates in Yangambi center.

According to destination of harvests (Table 2), most crop production is for sale with a smaller proportion for self-consumption (45.6% of respondents). Farmers who take all their harvests to market and keep nothing for self-consumption total 30.4%; 17.2% consume most of the harvest and sell a small portion; 3.2% divide the harvest equally between self-consumption and sale; and 3.6% practice self-consumption only. Fisher’s exact test showed that there is no significant difference in terms of crop destination between the axes ($p = 0.542$). However, a high proportion of farmers practicing agriculture for self-consumption only are found in Yangambi center.

During agricultural activities, farmers face many constraints. Hard work (reported by 40.9% of respondents), lack of means to pay for labor (18.6%), lack of road infrastructure (16.4%) and lack of farming equipment (16.4%) are the main constraints faced by our respondents. The lack of means of conservation and buyers were also mentioned.

Main crops and destination of harvests	Kisangani–Yangambi axis (<i>n</i> = 65)	Yangambi–Ngazi axis (<i>n</i> = 45)	Yangambi center (<i>n</i> = 140)	All Yangambi landscape (<i>n</i> = 250)	<i>p</i> -value
<i>Main crop</i>					0.004
Groundnut	2 (3.1)	5 (11.1)	6 (4.3)	13 (5.2)	
Plantain and table banana	2 (3.1)		3 (2.1)	5 (2.0)	
Maize	6 (9.2)	3 (6.7)	6 (4.3)	15 (6.0)	
Cassava	49 (75.4)	20 (44.4)	86 (61.4)	155 (62.0)	
Rice	6 (9.2)	17 (37.8)	33 (23.6)	56 (22.4)	
Cowpea			6 (4.3)	6 (2.4)	
<i>Destination of harvests</i>					0.542
Self-consumption	2 (3.1)		7 (5.0)	9 (3.6)	
Market_selfcons	4 (6.2)		4 (2.9)	8 (3.2)	
Market	21 (32.3)	14 (31.1)	41 (29.3)	76 (30.2)	
Plus_selfcons	9 (13.8)	10 (22.2)	24 (17.1)	43 (17.2)	
Plus_Market	29 (44.6)	21 (46.7)	64 (45.7)	114 (45.6)	

Table 2. Main crops and destination of harvests, expressed as: number of respondents (percentage of respondents)

Notes: Legend: Plus_Market = most of the harvest is destined for sale, Market = all the harvest is destined for sale, Plus_selfcons = most of the harvest is destined for self-consumption, Market_selfcons = the proportion sold is equal to the proportion self-consumed, Self-consumption = all the harvest is for self-consumption, X(Y): X = frequency and Y = percentage

3.2 Farmers' perceptions of climate change in the Yangambi landscape

3.2.1 Perceptions about climate change. The vast majority (98 %) of respondents perceived changes in temperature, rainfall and weather patterns (Table 3). Thus, 52.4% reported an increase in temperature; 45.2% reported unpredictable rainfall; 27.6% reported a decrease in the amount of rainfall per year; and 27.2% reported an increase in the amount of rainfall per year. Onset and cessation of rainfall creating disruptions to the agricultural calendar (52.4%), strong wind (21.6%) and delayed rainfall (7.2%) were also reported.

Fisher's exact test showed that observation of climate variations does not depend on the axes ($p = 0.244$). This is independent of gender ($p = 0.192$), age ($p = 0.387$), level of education ($p = 0.432$), farmers' main activity ($p = 0.343$) and farmers' seniority in agricultural activities ($p = 0.565$). However, there is a significant difference between axes in terms of decrease and increase in rainfall per year ($p < 0.001$). The decrease is observed more often along the Kisangani–Yangambi axis, whereas the increase is most reported along the Yangambi–Ngazi axis.

Climate variations mentioned above have been observed for 4.5 ± 6.1 years in the Yangambi landscape, with a minimum of 1 year and maximum of 36 years (data not shown). Regarding axes, observations have been made on average along the Kisangani–Yangambi axis for 6.3 years, 4.6 years in Yangambi center and 1.8 years along the Yangambi–Ngazi axis. Thus, it appears that proximity to the large city of Kisangani has an impact on climate variations, the closest sites being the most disturbed.

Farmers attribute observed climate variations to different causes. Although 47.6% do not know the real cause of the changes, 32% attribute them to divine phenomena (God's will/punishment being fulfilled), 7.2% to deforestation, 4.8% to demography and 3.6% to slash-and-burn agriculture. Fisher's exact test showed that knowledge about the causes of the changes depends on the level of education ($p = 0.002$), the seniority of the farmer in agricultural activities ($p = 0.019$) and the main activity of each farmer ($p < 0.001$). Farmers with higher education, i.e. university, cited slash-and-burn agriculture, whereas those with a secondary school diploma mentioned deforestation, demography and slash-and-burn agriculture. Those with a primary school level and those without education alluded to divine phenomena.

Perceptions of respondents	Axes			All Yangambi landscape	<i>p</i> -value
Perception based on climate change indicators	Kisangani–Yangambi axis (<i>n</i> = 65)	Yangambi–Ngazi axis (<i>n</i> = 45)	Yangambi center (<i>n</i> = 140)	All Yangambi landscape (<i>n</i> = 250)	
<i>Observation of Climate change</i>					
<i>Temperature trends</i>					
Increasing	62 (95.4)	45 (100.0)	138 (98.6)	245 (98)	0.244
<i>Rainfall trends</i>					0.041
Increasing	35 (53.8)	16 (35.6)	80 (57.1)	131 (52.4)	
Decreasing	6 (9.23)	20 (44.4)	42 (30.0)	68 (27.2)	<0.001
Unpredictable	32 (49.23)	6 (13.3)	31 (22.1)	69 (27.6)	
<i>Weather changes experienced</i>					
Delayed rainfall	27 (41.53)	19 (42)	67 (47.9)	113 (45.2)	
Early rainfall	2 (3.1)	6 (13.3)	10 (7.1)	18 (7.2)	0.129
Strong wind	0 (0)	2 (4.4)	3 (2.1)	5 (2)	0.301
Onset and cessation of rainfall	10 (15.4)	12 (26.7)	32 (22.9)	54 (21.6)	0.309
Other changes	32 (49.2)	27 (60.0)	72 (51.4)	131 (52.4)	0.528
	4 (6.2)	0 (0)	3 (2.1)	7 (2.8)	0.168

Notes: Legend: X(Y): X = frequency; Y = percentage

Table 3. Climate change perception in the yangambi landscape based on climate change indicators, expressed as: number of respondents (percentage of respondents)

3.2.2 Impacts of climate change on agricultural activities. Climate variations in the Yangambi landscape impact negatively agricultural yields/production. Indeed, 54.8% of respondents reported reduced crop yields; 43.6% reported the appearance of various new crop pests; 69.6% reported the appearance of new weed species; and 22.4% reported crop wilt (Table 4). The most affected crops reported by respondents are rice (31.2%), cowpea (20 %) and maize (18 %). Rice, maize and cowpea are most affected by the increase in temperature and decrease in rainfall, whereas cassava and groundnut are most affected by the increase in rainfall.

According to axes, Fisher's exact test showed that there is a significant difference in observation of yield reduction ($p < 0.001$) and crop new pest occurrence ($p = 0.001$) between axes. Yield reduction is most marked in Yangambi center (reported by 66.4% of respondents), followed by the Yangambi–Ngazi axis (44.4%) and the Kisangani–Yangambi axis (36.9%). The appearance of crop pests is more pronounced along the Kisangani–Yangambi axis. The appearance of new weed species has been observed on average for 8.3 ± 7.8 years, with a minimum of 1 year and maximum of 38 years. New pests have been observed on average for 3.3 ± 2.0 years, with a minimum of 1 year and maximum of 10 years.

3.2.3 Crop pests and weeds emerging as a result of climate variations. The most cited new weed species are *Croton hirtus* L'HÉRIT (reported by 52.4% of respondents), commonly called "abisibisi", a word from the Turumbu local language "abisi" meaning "to tickle"; *Mitracarpus villosus* (SW.) DC. (21.2% of respondents), locally called "botola", a word from the Lingala local language, "kobotola" meaning "to delight"; and *Oldenlandia corymbosa* L. (16.8% of respondents), locally called "ebola", in reference to the "Ebola virus" disease because of the damage it causes. Others weeds such as *Chromolaena odorata* (L.) R. King and H. Rob (14.9% of respondents) and *Hypparrhenia familiaris* (Steud.) Stapf were also mentioned (Table 5).

Impacts on agricultural Activities	Impacts of climate variability on crop production/yield	Axes			All Yangambi landscape	<i>p</i> -value
		Kisangani–Yangambi axis (<i>n</i> = 65)	Yangambi–Ngazi axis (<i>n</i> = 45)	Yangambi center (<i>n</i> = 140)		
Reduction of crops yield	24 (36.9)	20 (44.4)	93 (66.4)	137 (54.8)	<0.001	
Appearance of crop pests	41 (63.1)	15 (33.3)	53 (37.9)	109 (43.6)	0.001	
Appearance of new weed species	51 (78.5)	32 (71.1)	91 (65.0)	174 (69.6)	0.146	
Crop wilt	10 (15.4)	5 (11.1)	41 (29.3)	56 (22.4)	0.012	
Others	4 (6.2)	1 (2.2)	3 (2.1)	8 (3.2)	0.331	
<i>Most affected crops</i>						
Rice	10 (15.4)	9 (20.0)	59 (42.1)	78 (31.2)	<0.001	
Maize	26 (40.0)	7 (15.6)	12 (8.6)	45 (18)	<0.001	
Cowpea	5 (7.7)	4 (8.9)	41 (29.3)	50 (20)	<0.001	
Cassava	5 (7.7)	3 (6.7)	24 (17.1)	32 (12.8)	0.079	
Groundnut	5 (7.7)	5 (11.1)	14 (10.0)	24 (9.6)	0.839	
Other	6 (9.2)	1 (2.2)	10 (7.1)	17 (6.8)	0.360	

Notes: Legend: X(Y): X = frequency; Y = percentage

Table 4. Impacts of climate change on agricultural activities in the Yangambi landscape, expressed as: number of respondents (percentage of respondents)

Climate impacts	Weed/pest species	Kisangani–Yangambi axis	Yangambi–Ngazi axis	Yangambi center	All Yangambi landscape	<i>p</i> -value	
		(<i>n</i> = 65)	(<i>n</i> = 45)	(<i>n</i> = 140)	(<i>n</i> = 250)		
New weed species	<i>Croton hirtus</i>	42 (64.6)	26 (57.8)	63 (45.0)	131 (52.4)	0.024	
	<i>Pteridium aquilinum</i>		2 (4.4)	13 (9.3)	15 (6)	0.017	
	<i>Mitracarpus villosus</i>		17 (37.8)	36 (25.7)	53 (21.2)	<0.001	
	<i>Bulbostylis hispidula</i>	9 (13.8)	2 (4.4)		11 (4.4)	<0.001	
	<i>Chromolaena odorata</i>	23 (35.4)		3 (2.1)	26 (10.4)	<0.001	
	<i>Oldenlandia corymbosa</i>			42 (30.0)	42 (16.8)	<0.001	
	Others	8 (12.3)	7 (15.6)	11 (7.9)	26 (10.4)	0.249	
	New crop pests	<i>Zonocerus variegatus</i>	38 (58.5)	13 (28.9)	31 (22.1)	82 (32.8)	<0.001
		<i>Spodoptera frugiperda</i>	9 (13.8)	2 (4.4)	4 (2.9)	15 (6)	0.011
		<i>Aphis craccivora</i>	2 (3.1)	1 (2.2)	22 (15.7)	25 (10)	0.002
Others		5 (7.7)		2 (1.4)	7 (2.8)	0.028	

Notes: Legend: X(Y): X = frequency; Y = percentage

Table 5. Crop pests and weeds resulting from climate variations, expressed as: number of respondents (percentage of respondents)

However, it should be noted that for farmers, not all these weeds have negative effects. This is the case of *Chromolaena odorata*, locally called “likanja ikondo” in the Turumbu local language, which means “foreign”. Farmers prefer it to grow after harvest, thus contributing to the restoration of soil fertility in fallow land.

Regarding new pests that have emerged in recent years, 32.8% of farmers cited the variegated grasshopper *Zonocerus variegatus* (Linnaeus, 1758) (locally called “kwalala”);

10% reported the aphid *Aphis craccivora* Koch, 1854; and 6% reported the fall armyworm *Spodoptera frugiperda* (J. E. Smith, 1797).

3.3 Adaptation and resilience to climate change

In the case of yield reduction, most respondents have no adaptation techniques (87.6% of respondents). A small proportion adapt by directly resuming the seeding of the same crop (5.6% of respondents), by crops rotation (2.8% of respondents) or by clearing a new field in the forest (2% of respondents). Fisher’s exact test showed that adaptation techniques depend on the axis ($p = 0.027$) and on the farmer’s main activity ($p < 0.001$). Adaptation is independent of the level of education, age, membership of a peasant association and the farmer’s seniority in agricultural activities. Farmers in Yangambi center have a greater diversity of possibilities to adapt than do other farmers.

However, even if they are adopted by a small proportion of respondents, some practices used as solutions can be considered as CSA practices, including use of fertilizers, back to fallow, crops rotation and bio-pesticides. Table 6 below presents the adaptation/resilience options used, sorted as CSA and non-CSA practices in order to show what CSA option can be easily promoted/implemented in the region. Note that CSA practices are those that offer the triple possibility of simultaneously raising productivity, enhancing adaption/resilience and mitigating carbon emissions.

When new crop pests appear, 72.5% of respondents have no means of control, while 22% use mechanical control and 4.5% use bio-pesticides. Of the three axes, pesticides are used only in Yangambi center. This shows that vulnerability to pests increases with increased distance from the INERA/Yangambi national research center.

Apart from the abovementioned options already used by farmers (data not shown), there are other options that they believe to be effective even if they do not yet use them. The proposed options are scientific research (reported by 43.2% of respondents), recourse to God (25 % of respondents) and reforestation (9% of respondents). A proportion of 2.3% of respondents reported that agricultural activities should be reduced and other activities should be considered.

Practice reported	Kisangani– Yangambi axis (n = 65)	Axes Yangambi– Ngazi axis (n = 45)	Yangambi center (n = 140)	All Yangambi landscape All Yangambi landscape (n = 250)	CSA p-value practice
<i>Agricultural yields reduction</i>					0.027
No adaptation tracks	65 (100.0)	40 (88.9)	114 (81.4)	219 (87.6)	
Directly resume seeding		5 (11.1)	9 (6.4)	14 (5.6)	–
Clearing a new field in forest			5 (3.6)	5 (2.0)	–
Fertilizers			1 (0.7)	1 (0.4)	+
Back to fallow			2 (1.4)	2 (0.8)	+
Crops rotation			7 (5.0)	7 (2.8)	+
Other			2 (1.4)	2 (0.8)	
<i>Crops pests</i>					0.006
No adaptation tracks	49 (75.4)	40 (88.9)	118 (84.3)	207 (82.8)	
Bio-pesticides			9 (6.4)	9 (6.4)	+
Mechanical control	15 (23.1)	5 (11.1)	13 (9.3)	33 (13.2)	–

Notes: Legend: + = CSA practice, – = non-CSA practice, X(Y): X = frequency and Y = percentage

Table 6.
Adaptation and resilience to climate change, expressed as: number of respondents (percentage of respondents)

In addition, to promote agricultural activities in the Yangambi landscape, a large proportion of respondents need farming equipment and need the advice of an appropriate agricultural service to cope with these new challenges. Good agricultural feeder roads, fertilizers, pesticides and improved varieties were also reported.

3.4 Climate-smart agriculture framework

According to respondents' perceptions, [Table 7](#) proposes a CSA framework for the Yangambi landscape. It lists most reported climate risks, their impacts on agricultural activities and the likely pathways for CSA options that have been adapted to the realities of the Yangambi landscape.

4. Discussion

4.1 Agricultural practices

In the Yangambi landscape, each household owns 1.67 ± 0.79 fields of 0.75 ± 0.52 ha, mainly established on fallow land. These results are close to those found by [Kyale et al. \(2019\)](#) in the same region. Field sizes are close to the Tshopo Province average ([FONAREDD, 2016](#)) but small compared with the national average ([Potapov et al., 2012](#)).

The small field area can be explained by the presence of the Yangambi Biosphere Reserve. As farmers are not allowed to open new fields in the forest belonging to the protected area, their children are obliged to divide up the land that belonged to their parent to establish fields. This reality was noted by [Kyale and Maindo \(2017\)](#), who characterized access to land in this area as "essentially hereditary". This mode of access to land would also explain the abundance of fields established on fallow land. The same reality is observed in Gbere village located in the east of Garamba National Park ([Semeki et al., 2016](#)) and in villages bordering the Salonga National Park in DRC ([ICCN, 2018](#)).

The main crops in Yangambi landscape include cassava, rice, maize, groundnuts and plantain and table banana, which are also the main crops of Tshopo Province ([Moonen, 2017](#)). Our findings are similar to those of [Semeki et al. \(2016\)](#) in the northeast of the country and of the [ICCN \(2018\)](#) in the central part of the country. Cassava is the country's main crop ([Molinario et al., 2015](#)).

Most of the crop harvest is brought to market, as agriculture is subsistence farming ([Semeki et al., 2016](#)). The money obtained from the sale is used to pay children's school fees, buy farm equipment, pay debts and buy other nonagricultural products. In addition, a high proportion of farmers practicing agriculture for self-consumption only are found in Yangambi center. This can be explained by the fact that activities are more diversified in Yangambi center than in other areas.

The tree species found in respondents' fields are of some interest to farmers ([Batsi et al., 2020](#)). *Petersianthus macrocarpus* and *Persea americana* are the two preferred species; the former is home to edible caterpillars and the latter provides edible fruits. *P. macrocarpus* trees have been preserved in the Yangambi landscape because of the caterpillars ([Kyale and Maindo, 2017](#)).

4.2 Farmers' perceptions of climate change

4.2.1 Climate change perceptions. Climate variations are observed by the vast majority of respondents. The climate risks differ locally, nationally and regionally ([Ministry of Foreign Affairs, 2018](#)).

Climate variations have been mentioned by other researchers in the same region ([CIFOR, 2018b](#); [Likoko et al., 2019](#)) and in other regions of the country ([Bele et al., 2014](#); [Ulimwengu](#)

Climate risks	Impacts	CSA options	Potential partners/actors
Increase in temperature	Crop wilt	Use of tolerant varieties, mulching, agroforestry	Research institutions, state technical services, nongovernmental organizations (NGOs)
	Reduced crop yields	Use of tolerant varieties, agricultural support service, agroforestry, revival of agricultural technical schools, livelihood diversification and new crops, mulching	
	Delayed crop growth	Use of short-cycle varieties, fertilizer and manure application, mulching	
Increase in annual rainfall	Removal of pests	Use of bio-pesticides, consumption of variegated grasshoppers	Local community, research institutions, NGOs
	Removal of pests	Intensive weeding, high seeding density, livelihood diversification, new crops	
Decrease in annual rainfall	Cassava tuber rot	Use of tolerant varieties	State technical services, NGOs
	Crop wilt	Irrigation, water recycling, lowland development	
Onset and cessation of rainfall	Growth delay	Irrigation, use of resistant varieties, livelihood diversification, new crops	Research institutions, state and NGO technical services
	Erosion of rain-fed soils supporting rice, groundnuts, and maize	Use of tolerant varieties, change of planting date	
Strong winds	Cassava tuber rot	Use of improved seeds	Local communities, NGOs
	Agricultural yield reduction	Agricultural calendar adjustment, diversification of activities, livelihood diversification, new crops, early-maturing varieties, mulching	
Strong winds	Pruning multipurpose trees	Windbreaks, agroforestry, tree planting and protection	Local communities, NGOs
	Pouring of banana trees and other crops	Windbreaks, assisted natural regeneration, reforestation of degraded areas, agroforestry	

Table 7.
Climate-smart
agriculture
framework according
to the Yangambi
landscape realities

and Kibonge, 2016). They are also observed in all African regions (Sonwa *et al.*, 2016; Tesfahun and Chawla, 2019; Saalu *et al.*, 2020).

The increase in temperature, onset and cessation of rainfall, decreased rainfall in some areas and increased annual rainfall in other areas are the main climate risks observed in the Yangambi landscape. Likoko *et al.* (2019) mentioned an increase in temperature of 0.44°C between 1931 and 2017 in the same area. Findings of our study corroborate those reported by the National Capacity Needs Assessment (MECNT and UNDP, 2009) and Chinedum *et al.*

(2015) in DRC. They match those found by Chia *et al.* (2016) in Cameroon, Onyeneke *et al.* (2017) in Nigeria, Saalu *et al.* (2020) in Kenya and Oriangi *et al.* (2020) in Uganda.

Regarding the duration of climate variations, hydrometric and meteorological data of 17 national meteorological stations showed that variations have been observed since the 1980s throughout the DRC (MECNT and UNDP, 2009). In the Yangambi landscape, climate variations have been observed for an average duration of 4.5 ± 6.1 years, with a minimum of 1 year and maximum of 38 years. This maximum is close to the 37 years reported by Codjo *et al.* (2015) in Benin.

Even if a large proportion of respondents do not know the causes of these changes, some of them attribute it to divine phenomena (God's will/punishment being fulfilled), deforestation, slash-and-burn agriculture and demography. These findings match those found by Yila and Resurreccion (2014) in Nigeria and Tesfahun and Chawla (2019) in Ethiopia. However, slash-and-burn agriculture as a cause of climate change has been cited at local (Hufkens *et al.*, 2020), national (MECNT, 2012; Molinario *et al.*, 2015) and international levels (Pan *et al.*, 2011; Arneth *et al.*, 2019).

4.2.2 Impacts of climate change on agricultural activities. Impacts of climate variations are inevitable, and even in the most optimistic scenario, they will have adverse effects on agricultural production and yields (Brown *et al.*, 2011). They differ by country and region, as well as by community and individual (Ministry of Foreign Affairs, 2018).

Regional climate models of the Congo Basin predict that a combination of factors, i.e. increase in temperature, rainfall decrease and irregularity or even extreme situations will inevitably result in a significant reduction in agricultural production and yield, a greater sensitivity of crops to pest attacks and a decrease in soil fertility (Lo and Kaéré, 2009). Moiroux *et al.* (2014) and Cilas *et al.* (2015) pointed out that climate change leads to the proliferation of pests transferring from one region to another. Our findings show that the main impacts are the appearance of new weed species, crop yield reduction and the appearance of crop pests, which matches the above viewpoints. Particular emphasis has been placed on yield reduction in rain-fed farming by Bele *et al.* (2014) in eastern DRC, Doukpolo (2014) in Central Africa and Ulimwengu and Kibonge (2016) in western DRC.

Hailu and Campbell (2015) pointed out that cereals are the most vulnerable crops (especially rice and maize). This viewpoint is confirmed by our findings. In addition, our results match the findings of Saalu *et al.* (2020) in Kenya and those of the IPCC in many lower latitude regions (Arneth *et al.*, 2019).

4.2.3 Emerging crop pests and weeds. Peltier *et al.* (2014) reported that forests were being replaced by weeds such as *Chromolaena odorata*, *Pteridium aquilinum* and *Imperata cylindrica* around the city of Kinshasa. However, in the Yangambi landscape, these weeds are rarely mentioned, with the most cited weeds being *Croton hirtus*, *Mitracarpus villosus* and *Oldenlandia corymbosa*.

The variegated grasshopper *Zonocerus variegatus*, the aphid *Aphis craccivora* and the fall armyworm *Spodoptera frugiperda* are the most widespread new pests in the Yangambi landscape. Bakondongama *et al.* (2017) reported *Z. variegatus* in Tshopo Province as a real threat to agriculture. In addition, Goergen *et al.* (2016) noted that in 2016, a new caterpillar (*Spodoptera frugiperda*) arrived in Africa. The FAO mentioned it first in West Africa and Central Africa in early 2016; and in 2018, it was detected and reported in almost all sub-Saharan African countries (FAO, 2018).

4.3 Other constraints on agricultural activities

Apart from climate change-related constraints, agricultural activities and crop yields face other constraints. Hard work, lack of means to pay for labor, lack of road infrastructure and

lack of adequate farming equipment are the main constraints faced by our respondents. In 2018, [CIFOR \(2018b\)](#) reported in the same area, the lack of availability of agricultural inputs, lack of supervision of farmers and nonexistence of cooperatives ensuring the purchase and sale of agricultural products. At provincial level, [Van Hoof \(2011\)](#) pointed out that roads are in poor condition, the many administrative and police hassles and lack of quality seeds.

In the Congo Basin (Cameroon, Central African Republic and DRC), constraints on farmers that make their adaptive capacity weak are the lack of technology, information, skills and infrastructure. Specifically, these are, in Cameroon, the lack of infrastructure and human and financial resources ([Brown et al., 2011](#)); in Central African Republic, the lack of equipment, financial and technical resources and above all access to information ([Doukpolo, 2014](#)); and in DRC, the lack of basic infrastructure, producer organization, financial resources and processing activities ([MINAGRI, 2010](#)).

4.4 Possible solutions

Farmers do not stand idly by in the face of the negative impacts of climate change on agricultural activities. They are proposing certain solutions, either concrete or difficult to implement. Rural communities are the most vulnerable due to their limited adaptation capacities and their high dependence on climate-sensitive resources (IPCC, 2007; [Brown et al., 2011](#); [Bele et al., 2014](#)).

In the case of yield reduction, most farmers do not have adaptation/resilient techniques, while some use crops rotation, fallow practice, or replant the same type seed. However, others open up new fields in the forest, posing an additional risk to the forest by accelerating the climate change process through deforestation. These results show that adaptive capacity in the Yangambi landscape is low compared with farmers in other African countries, including Nigeria ([Onyeneke et al., 2017](#)) and Ethiopia ([Tesfahun and Chawla, 2019](#)). In terms of axes, farmers in Yangambi center have a diversity of solutions compared with other farmers. This may be due to their proximity to the National Institute for Agronomic Studies and Research, INERA/Yangambi.

[Chinedum et al. \(2015\)](#) reported eight CSA practices used by farmers at the national level in DRC. Our findings match some of them, especially as regards fertilizers, resumption of use of fallows, crops rotation and use of bio-pesticides. However, these practices are adopted by a very small proportion of respondents and are not yet being promoted. Note that agroforestry is already practiced by farmers in some corners of the Yangambi landscape ([Batsi et al., 2020](#)) but only by a few. [Onyeneke et al. \(2017\)](#) reported that farmers have already adopted CSA practices in southeast Nigeria.

At the national level, to combat climate change, the DRC has set up the PANA program. Its objective is to develop a program covering the entire country and identifying urgent and immediate adaptation activities that respond to the current and anticipated adverse effects of climate change ([PANA, 2006](#)). This program presents agriculture as the sector most vulnerable to climate change in DRC ([MINAGRI, 2010](#)). The main component of this project concerns the revival of the vital sector of basic agricultural production with a view to achieving food security. The project was carried out in four intervention zones located in the provinces of Katanga, Kasai Oriental, Bandundu and Bas-Congo ([PANA, 2006](#)). Even though it was recommended, in terms of future directions, that particular attention must be paid to carrying out in-depth studies of climatic, socio-economic, cultural and ecological vulnerability to prepare zone-specific responses; to date, nothing has yet been done in the Yangambi landscape under PANA.

In 2008, DRC benefited from the Congo Basin Forests and Adaptation to Climate Change (CoFCCA) project. Its objective was to assess the level of vulnerability to climate change of

local communities in the forests of the Congo Basin and to bring stakeholders, through the participatory action research approach, to design and implement specific adaptation strategies (Sonwa *et al.*, 2014). The study carried out using this approach revealed that the adaptation efforts developed by local communities are no longer sufficient in the face of climatic uncertainties and the speed at which these changes are occurring (Bele *et al.*, 2010).

In the same framework as the fight against climate change, a mechanism called REDD+, of which the DRC has been part since 2009, was launched at the global level. The objectives of the national REDD+ strategy in DRC are to contribute to the mitigation of GHG emissions, poverty reduction and sustainable management of its forest resources, taking into account the valuation of environmental services (MECNT, 2015). The DRC's REDD+ program launched an eastern integrated program in 2016. It aimed at:

- improvement of governance of natural resources; and
- reduction of the impact of economic activities and population dynamics on forests.

However, it should be noted that the Yangambi landscape was not part of the project area. Agriculture was considered for the sole purpose of reducing deforestation (FONAREDD, 2016).

From previous activities, it appears that projects under REDD+ and the national PANA program have not specifically addressed the Yangambi landscape. In addition, agriculture is generally involved by mainly targeting a few main objectives (food security, climate mitigation, climate adaptation or biodiversity conservation) over a specific time period and geographic locations (Sonwa *et al.*, 2020).

Thus, the promotion of agriculture through CSA techniques is crucial for the Yangambi landscape and the well-being of the local population, as it links three important objectives according to the local context of climate change. Its implementation needs support and coordination from many stakeholders around smallholder farmers, including from research institutions, nongovernmental organizations (NGOs) and government institutions (Bele *et al.*, 2010; Sonwa *et al.*, 2020).

The African Ministerial Conference on the Environment (AMCEN, 2014) mentioned that science is the basis for decision-making for Africa, whenever the challenge of climate change needs to be understood or solutions need to be developed. African universities could become centers of innovation and technology transfer for CSA (Hailu and Campbell, 2015). In addition, government policies need to support research and development that focuses on and diffuses climate-smart technologies to help farmers respond to climate variations (Onyeneke *et al.*, 2017).

5. Conclusion

In the Yangambi landscape, farmers perceive climate variations in temperature, rainfall and weather patterns. Increase in temperature, onset and cessation of rainfall creating disruptions in the agricultural calendar and decrease or increase in annual rainfall are the main climate risks. Their impacts on agricultural activities are that many crop yields are reduced, and that new weeds species and new crop pests emerge.

Although a large proportion of respondents have no adaptation and resilience techniques, some of them use crops rotation, fallow practice, fertilizers and bio-pesticides, which can be considered as CSA practices already used in the region. This indicates that even if farmers are not yet receiving external assistance from government or non-governmental organizations to adapt and cope with climate change, they are able to put some CSA techniques into practice. To do so, it is therefore necessary to valorize these

practices and popularize them among farmers in the Yangambi Landscape, as they are accessible and feasible at the local level.

On the other hand, CSA practices to be proposed in the Yangambi landscape according to local realities are multiple, including the adjustment of the agricultural calendar, agroforestry, the use of tolerant and improved varieties, crops rotation, development of lowlands and the use of bio-fertilizers and bio-pesticides. To promote the implementation of CSA in the Yangambi landscape, support from research institutions, NGOs and state institutions is necessary.

References

- Alongo, S., Visser, M., Drouet, T., Kombele, F., Colinet, G. and Bogaert, J. (2013), "Effets de la fragmentation des forêts par l'agriculture itinérante sur la dégradation de quelques propriétés physiques d'un ferralsol échantillonné à yangambi, R.D. Congo", *Tropicultura*, Vol. 31 No. 1, pp. 36-43.
- AMCEN (2014), "Strategie africaine sur les changements climatiques", AMCEN-15-REF-11, Addis-Ababa.
- Arnell, A., Barbosa, H., Benton, T., Calvin, K., Calvo, E., Connors, S., Cowie, A., Davin, E., Denton, F., Van Diemen, R., Driouech, F., Elbehri, A., Evans, J., Ferrat, M., Harold, J., Haughey, E., Herrero, M., House, J., Howden, M., Hurlbert, M., Jia, G., Johansen, T., Krishnaswamy, J., Kurz, Z., Lennard, C., Myeong, S., Mahmoud, N., Masson-Delmotte, V., Mbow, C., McElwee, P., Mirzabaev, A., Morelli, A., Moufouma-Okia, A., Nedjraoui, D., Neogi, S., Nkem, J., De Noblet-Ducoudré, N., Olsson, L., Pathak, M., Petzold, J., Pichs-Madruga, R., Poloczanska, E., Popp, A., Pörtner, H.-O., Pereira, J., Pradhan, P., Reisinger, A., Roberts, D., Rosenzweig, C., Rounsevell, M., Shevliakova, E., Shukla, P., Skea, J., Slade, R., Smith, P., Sokona, Y., Sonwa, D., Soussana, J.-F., Tubiello, F., Verchot, L., Warner, K., Weyer, N., Wu, J., Yassaa, N., Zhai, P. and Zommers, Z. (2019), "IPCC special issue on land degradation: Summary for policymakers", in Shukla, P., Skea, J., Buendia, E., Masson-Delmotte, V., Pörtner, H.-O., Roberts, D., Zhai, P., Slade, R., Connors, S., Van Diemen, R., Ferrat, M., Haughey, E., Luz, S., Neogi, S., Pathak, M., Petzold, J., Pereira, J., Vyas, P., Huntley, E., Kissick, K., Belkacemi, M. and Malley, J. (Eds.), *Climate Change and Land: An IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*, The Intergovernmental Panel on Climate Change (IPCC), Geneva. pp. 1-41.
- Bakondongama, B., Mubenga, K., Mondivudri, A., Ngabu, C., Mpiana, P., Kosele, K. and Juakaly, M. (2017), "Zonocerus variegatus (linné 1758, orthoptera: Pyrgomorphidae) et ses parasitoïdes (hyménoptères) dans la ville de kisangani, RDC", *International Journal of Innovation and Scientific Research*, Vol. 32 No. 1, pp. 64-74.
- Barnier, J., Biaudet, J., Briatte, F., Bouchet-Valat, M., Gallic, E., Giraud, F., Gombin, J., Kauffmann, M., Lalanne, C., Larmarange, J. and Robette, N. (2020), "Analyse-R – introduction à l'analyse d'enquêtes", *Avec R et RStudio*, doi: [10.5281/zenodo.2669067](https://doi.org/10.5281/zenodo.2669067), (accessed 15 December 2020).
- Batsi, G., Sonwa, D., Mangaza, L., Ebuy, J. and Kahindo, J.-M. (2020), "Biodiversity of the cocoa agroforests of the Bengamisa-Yangambi Forest landscape in the democratic republic of the Congo (DRC)", *Forests*, Vol. 11 No. 10, pp. 1-21.
- Bele, M., Sonwa, D. and Tiani, A.-M. (2014), "Local communities vulnerability to climate change and adaptation strategies in Bukavu in DR Congo", *The Journal of Environment and Development*, Vol. 23 No. 3, pp. 331-357.
- Bele, Y., Mulotha, E., Bokoto, B., Sonwa, D. and Tiani, A.-M. (2010), "Les effets du changement climatique dans le bassin du Congo: la nécessité de soutenir les capacités adaptatives locales", *Aperçu Sur L'adaptation*, Vol. 3, pp. 1-3.
- Brown, C., Nkem, J., Sonwa, D. and Bele, Y. (2011), *Capacité D'adaptation Institutionnelle et Réponse Au Changement Climatique Dans Les Forêts Du Bassin Du Congo Au Cameroun*, CIFOR, Bogor.

- Chia, E., Tiani, A.-M., Sonwa, D., Perez-Teran, A. and Tchatchou, B. (2016), "Securing well-being with the advent of climate hazards: Case of Forest-dependent communities in a landscape in the Congo basin", *International Journal of Climate Change Strategies and Management*, Vol. 8 No. 2, pp. 175-193.
- Chinedum, N., Tambi, E. and Bangali, S. (2015), *State of Knowledge on CSA in Africa: Case Studies from Nigeria, Cameroun and the Democratic Republic of Congo*, Forum for Agricultural Research in Africa, Accra.
- CIFOR (2018a), *Agro-Écologie et Agroforesterie Pour Une Gestion Durable Des Forêts*, CIFOR, Bogor.
- CIFOR (2018b), *Libérer le Potentiel de L'agriculture: Analyse Des Chaines de Valeur Agricoles Dans le Paysage de Yangambi*, CIFOR, Bogor.
- Cilas, C., Goebel, F., Babin, R. and Avelino, J. (2015), "Bioagresseurs des cultures tropicales face au changement climatique: quelques exemples", in Torquebiau, E. (Ed.), *Changement Climatique et Agricultures Du Monde*, Quae, Versailles, pp. 75-83.
- Codjo, T., Ogouwale, E., Boko, M. and Agbossou, E. (2015), "Stratégies paysannes d'adaptation aux changements climatiques dans la commune d'Adjohoun", *XXVIII Colloque de L'Association Internationale de Climatologie, Liège 2015*, pp. 313-318.
- Delvaux, F. (2016), *L'agriculture Intelligente Face Au Climat, Les Fausses Réponses Au Changement Climatique*, Vol. 4, Entraide et Fraternité, Bruxelles.
- Doukpolo, B. (2014), "Changements climatiques ", et productions agricoles dans l'Ouest de la République Centrafricaine", PhD thesis, Université d'Abomey-Calavi.
- FAO (2014), *Exemples de Réussites de la FAO en Matière D'agriculture Intelligente Face au Climat*, FAO, Rome.
- FAO (2016), *Le Climat Change, L'alimentation et L'agriculture Aussi*, FAO, Rome.
- FAO (2018), *Gestion Intégrée de la Chenille Légionnaire D'automne Sur le Maïs: un Guide Pour Les Champs-Écoles Des Producteurs en Afrique*, FAO, Rome.
- FONAREDD (2016), *Fonds National REDD + de La RDC: Programme Intégré Oriental*, FONAREDD, Kinshasa.
- Goergen, G., Kumar, L., Sankung, S., Togola, A. and Tamo, M. (2016), "First report of outbreaks of the fall armyworm *Spodoptera frugiperda* (J E smith) (lepidoptera, noctuidae), a new alien invasive pest in West and Central Africa", *PLoS One*, Vol. 11 No. 10, pp. 1-9.
- Hailu, M. and Campbell, B. (2015), *L'agriculture Intelligente Face Au Climat: Quel Impact Pour L'Afrique? Success Stories*, CTA, Wageningen.
- Hufkens, K., De Haulleville, T., Kearsley, E., Jacobsen, K., Beeckman, H., Stoffelen, P., Vandeloek, P., Meeus, S., Amara, M., Van Hirtum, L., Van den Bulcke, J., Verbeeck, H. and Wingate, L. (2020), "Historical aerial surveys map long-term changes of Forest cover and structure in the Central Congo basin", *Remote Sensing*, Vol. 12 No. 4, pp. 1-26.
- ICCN (2018), *Rapport Sur L'état de Conservation Des Biens de la RDC Inscrits Sur la Liste du Patrimoine Mondial en Péril: exercice 2017*, ICCN, Kinshasa.
- IPCC (2007), "Climate change 2007: Synthesis report", *Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC)*, Geneva.
- IPCC (2014), "Climate change 2014: Synthesis report", *Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC)*, Geneva.
- Kipalu, P. and Mukungu, J. (2012), Statut Du Processus REDD + En République " Démocratique Du Congo", *Forest Peoples Programme*, Birmingham.
- Kombebe, B. (2004), "Diagnostic de la fertilité des sols dans la Cuvette Centrale congolaise. Cas des séries yakonde et yangambi", PhD thesis, Faculté Universitaire des Sciences Agronomiques de Gembloux, Gembloux.
- Kyale, J. and Maindo, A. (2017), "Pratiques traditionnelles de conservation de la nature à l'épreuve des faits chez les peuples riverains de la réserve de biosphère de yangambi (RDC)", *European Scientific Journal*, Vol. 13 No. 8, pp. 328-356.

- Kyale, J., Wardell, D., Mikwa, J.-F., Kabuanga, J., Maindo, A., Oszwald, J. and Doumenge, C. (2019), "Dynamique de la déforestation dans la réserve de biosphère de yangambi (république démocratique du Congo): variabilité spatiale et temporelle au cours des 30 dernières années", *Bois and Forêts Des Tropiques*, Vol. 341 No. 3, pp. 15-28.
- Likoko, A., Mbifo, N., Besango, L., Totiwe, T., Badjoko, H., Likoko, G., Botomo, D., Litemandia, N., Posho, B., Alongo, S. and Boyemba, F. (2019), "Climate change for Yangambi Forest region, DR Congo", *Journal of Aquatic Sciences and Oceanography*, Vol. 1 No. 2, pp. 1-10.
- Lo, H. and Kaéré, A. (2009), "L'Afrique et les changements climatiques: enjeux et défis multiples", in Mbow, A., Diop, S., Tounkara, A. and Lamine, M. (Eds.), *AGRIPADE: Revue Sur L'agriculture Durable à Faibles Apports Externes*, Vol. 24, AGRIPADE, Dakar, pp. 6-7.
- MECNT (2012), *Synthèse Des Études Sur Les Causes de La Déforestation et de La Dégradation Des Forêts En République Démocratique Du Congo*, MECNT, Kinshasa.
- MECNT (2015), *Rapport de Base: Analyse Environnementale et Sociale Stratégique Du Processus REDD+*, MECNT, Kinshasa.
- MECNT and UNDP (2009), *Autoévaluation Nationale Des Besoins En Renforcement de Capacités Pour La Gestion de L'Environnement En République Démocratique Du Congo*, MECNT, Kinshasa.
- MINAGRI (2010), *Programme REDD + Réduction de L'impact de L'agriculture de Subsistance Sur La Forêt: Document D'orientation*, MINAGRI, Kinshasa.
- Ministry of Foreign Affairs (2018), *Climate Change Profile: Democratic Republic of the Congo (East)*, Ministry of Foreign Affairs of the Netherlands, The Hague.
- Mohyont, B. and Demarée, G. (2006), "Courbes intensité-durée-fréquence des précipitations à yangambi, Congo, au moyen de différents modèles de type Montana", *Hydrological Sciences Journal*, Vol. 51 No. 2, pp. 239-253.
- Moiroux, J., Bourgeois, G., Boivin, G. and Brodeur, J. (2014), *Impact Différentiel Du Réchauffement Climatique Sur Les Insectes Ravageurs Des Cultures et Leurs Ennemis Naturels: Implications En Agriculture, Feuille Technique Ouranos Projet 550005-103*, Québec.
- Molinario, G., Hansen, M. and Potapov, P. (2015), "Forest cover dynamics of shifting cultivation in the democratic republic of Congo: a remote sensing-based assessment for 2000-2010", *Environmental Research Letters*, Vol. 10 No. 9, pp. 1-15.
- Mooney, P. (2017), "Integrated assessment of options to reduce emissions from deforestation in tshopo, DR Congo", PhD thesis, KU Leuven.
- Onyeneke, R., Igberi, C., Uwadoka, C. and Aligbe, J. (2017), "Status of climate-smart agriculture in southeast Nigeria", *GeoJournal*, Vol. 83 No. 2, pp. 333-346.
- Oriangi, G., Albrecht, F., Di Baldassarre, G., Bamutaze, Y., Mukwaya, P., Ardö, J. and Pilesjö, P. (2020), "Household resilience to climate change hazards in Uganda", *International Journal of Climate Change Strategies and Management*, Vol. 12 No. 1, pp. 59-73.
- Pan, Y., Birdsey, R., Fang, J., Houghton, R., Kauppi, P., Kurz, W., Phillips, O., Shvidenko, A., Lewis, S., Canadell, J., Ciais, P., Jackson, R., Pacala, S., McGuire, D., Piao, S., Rautiainen, A., Sitch, S. and Hayes, D. (2011), "A large and persistent carbon sink in the world's forests", *Science*, Vol. 333 No. 6045, pp. 988-993.
- PANA (2006), *Programme D'Action National D'Adaptation Au Changement Climatique de La République Démocratique Du Congo*, PANA, Kinshasa.
- Peltier, R., Dubiez, E., Diowo, S., Gigaud, M., Marien, J.-N., Marquant, B., Peroches, A., Proce, P. and Vermeulen, C. (2014), "Assisted natural regeneration in slash-and-burn agriculture: Results in the democratic republic of the Congo", *Bois and Forêts Des Tropiques*, Vol. 321 No. 321, pp. 67-79.
- Potapov, P., Turubanova, S., Hansen, M., Adusei, B., Broich, M., Altstatt, A., Mane, L. and Justice, C. (2012), "Quantifying Forest cover loss in democratic republic of the Congo, 2000-2010, with landsat ETM+ data", *Remote Sensing of Environment*, Vol. 122, pp. 106-116.

- Saalu, F., Oriaso, S. and Gyampoh, B. (2020), "Effects of a changing climate on livelihoods of Forest dependent communities: Evidence from buyangu community proximal to kakamega tropical rain Forest in Kenya", *International Journal of Climate Change Strategies and Management*, Vol. 12 No. 1, pp. 1-21.
- Seguin, B. (2010), "Le changement climatique: conséquences pour les végétaux", *Quaderni*, Vol. 71 No. 71, pp. 27-40.
- Semekki, J., Linchant, J., Quevauvillers, S., Kahindo, J.-M., Lejeune, P. and Vermeulen, C. (2016), "Cartographie de la dynamique de terroirs villageois à l'aide d'un drone dans les aires protégées de la république démocratique du Congo", *Bois and Forêts Des Tropiques*, Vol. 330 No. 330, pp. 69-83.
- Sonwa, D., Mangaza, L., Batsi, G. and Ebuy, J. (2020), "Preliminary reflections of the potential of climate smart agriculture (CSA) in the tshopo province (DRC)", in Matsunami, Y. and Takeuchi, S. (Eds.), *Challenges of Development and Natural Resource Governance in Africa; ASC–TUFS Working Papers 2019*, Tokyo, pp. 219-233.
- Sonwa, D., Nkem, J., Idinoba, M., Bele, M. and Jum, C. (2012), "Building regional priorities in forests for development and adaptation to climate change in the Congo basin", *Mitigation and Adaptation Strategies for Global Change*, Vol. 17 No. 4, pp. 441-450.
- Sonwa, D., Dieye, A., Mzouri, E., Majule, A., Mugabe, F., Omolo, F., Wouapi, H., Obando, J. and Brooks, N. (2016), "Drivers of climate risk in African agriculture", *Climate and Development*, Vol. 9 No. 5, pp. 383-398.
- Sonwa, D., Scholte, P., Pokman, W., Schauerte, P., Tsalefac, M., Biona, C., Brown, C., Haensler, A., Ludwig, F., Mkankam, F., Mosnier, A., Moufouma-Okia, W., Ngana, F. and Tiani, A. (2014), "Climate change and adaptation in centration africa: past, scenarios and options for the future", in De Wasseige, C., Flynn, J., Louppe, D., Hiol Hiol, F. and Mayaux, P. (Eds.), *The Forests of the Congo Basin-State of the Forest 2013*, Weyrich, pp. 99-119.
- Tesfahun, A. and Chawla, A. (2019), "Risk perceptions and adaptation strategies of smallholder farmers to climate change and variability in North shoa zone, Ethiopia", *Management of Environmental Quality: An International Journal*, Vol. 31 No. 1, pp. 254-272.
- Ulimwengu, J. and Kibonge, A. (2016), "Climate-smart agriculture practices based on precision agriculture: the case of maize in Western Congo", in De Pinto, A. and Ulimwengu, J.M. (Eds.), *A Thriving Agricultural Sector in a Changing Climate: Meeting Malabo Declaration Goals through Climate-Smart Agriculture*, USA, ReSAKSS Annual Trends and Outlook, Washington, DC DC, pp. 86-102.
- Van Hoof, F. (2011), *Changer L'agriculture Congolaise En Faveur Des Familles Paysannes*, AFAFO, Netherlands.
- Van Wambeke, A. and Evrard, C. (1957), *Notice Plicatice de La Carte Des Sols et de La Végétation. Planchette1*, INEAC, Bruxelles.
- Williams, T., Mul, M., Cofie, O., Kinyangi, J., Zougmore, R., Wamukoya, G., Nyamisi, M., Mapfumo, P., Ifejike, C., Amwata, D., Frid-Nielsen, S., Partey, S., Girvetz, E., Rosenstock, T. and Campbell, B. (2015), "Climate smart agriculture in the African context", paper presented at the high level conference on Feeding Africa: an action plan for African agriculture transformation, Senegal, 21–23 October 2015, Dakar.
- Yila, J. and Resurreccion, B. (2014), "Gender perspectives on agricultural adaptation to climate change in drought-prone nguru local government area in the semiarid zone of northeastern Nigeria", *International Journal of Climate Change Strategies and Management*, Vol. 6 No. 3, pp. 250-271.

About the authors

Lisette Mangaza is a Teaching Assistant and Researcher at the University of Goma (DRC), Faculty of Science, Department of Environment. She holds master's degree in biodiversity management and sustainable forestry planning (Forest and environment option) from the University of Kisangani

(DRC). Her research focuses on tropical forest ecology, climate change and sustainable agriculture. She is author and co-author of several scientific publications. Lisette Mangaza is the corresponding author and can be contacted at: lisemangaza@gmail.com

Denis Jean Sonwa is a Senior Scientist with CIFOR. His research activities are on Forest and Climate Change responses (Adaptation and Mitigation) in Central Africa. He previously worked with the International Institute of Tropical Agriculture on cocoa-agroforest management in West and Central Africa. Denis had been visiting professor at ASC-TUFS in Tokyo (Japan). He was trained at the University of Yaoundé (Cameroon), Defended PhD at the University of Bonn (Germany) and obtained a training certificate from the University of California, Berkeley (USA). He has authored and co-authored many publications including the recent IPCC special report on climate change and Land Degradation.

Germain Batsi is a Teaching Assistant and Researcher at the University of Kisangani, DRC. His research focuses on biodiversity conservation, climate change, deforestation and the improvement of peasant socio-economic conditions through the promotion of a sustainable agricultural system in Central Africa. He obtained master's degree in biodiversity management and sustainable forestry planning at the Faculty of Science of the University of Kisangani. He trained in Agricultural Engineering at the Faculty Institute of Agricultural Sciences of Yangambi. He is author and co-author of several scientific publications.

Jérôme Ebuy holds PhD in Forestry Sciences from the University of Kisangani. He attended the Université Catholique de Louvain (UCL) and the Royal Museum of Central Africa (RMCA) for his doctoral research. In addition, he has been trained in Sustainable Forest Management; REDD+ and Climate Change. His research area is Carbon Management (stocks and flows) under forest plantations (silviculture). Forestry graduate, he is currently Vice Dean in charge of research, Water and Forestry Unit, Faculty of Management of Renewable Natural Resources of the Université of Kisangani (DRC).

Jean-Marie Kahindo Muhongya holds Doctorate in Science from the University of Kisangani in DR Congo since 2011. A trained botanist, the Professor is interested in the interactions between man and the forest. He is the author and co-author of several scientific articles in the fields of Forest Ecology and Remote Sensing. Currently, he is responsible for the Chair of General Botany at the Department of Ecology and Plant Resource Management and Vice-President of the internship at the Faculty of Sciences, University of Kisangani, DRC.

For instructions on how to order reprints of this article, please visit our website:

www.emeraldgrouppublishing.com/licensing/reprints.htm

Or contact us for further details: permissions@emeraldinsight.com